








For the Infant and Child

							
Doses	Birth - ½ month (0 - 2 weeks)	½ - 1 ½ months (2 - 6 weeks)	1 ½ to 9 months (6 to 39 weeks)	9 - 12 months (39 – 52 weeks)	1- 2 years (12-24 months)	5 th year	10 & 16 years
OPV 0	Give in first 2 weeks only						
BCG	Give as early as possible in the first 12 months						
DPT 1 OPV 1	Give at 6 wks or as soon as possible after 6 wks						
DPT 2 OPV 2	Give at 10 wks/ as soon as possible after 10 wks <i>(wait at least 4 wks after DPT 1 & OPV 1 to give DPT 2 & OPV 2)</i>						
DPT 3 OPV 3	Give at 14 wks/as soon as possible after 14 wks <i>(wait at least 4 wks after DPT 2 & OPV 2 to give DPT 3 & OPV 3)</i>						
Measles	Give at 39 wks/ as soon as possible after 39 weeks						
DPT- OPV Booster	If a child does not receive Measles by 1yr, give as soon as possible before 5 yrs						
DT	Give at 16 months <i>(wait at least 6 months after OPV3 & DPT3)</i>						
TT	Give at 5 yrs/ school entry						
Vitamin A	Give at 10 and 16 years						
	Give at 9 th , 16 th 24 th 30 th 36 th 42 nd 48 th 54 th and 60 th month						